

## MAT-509

### Integral Transforms and Integral Equations

M. Sc. MATHEMATICS (MSCMAT-12)

Second Year, Examination, 2017

**Time : 3 Hours**

**Max. Marks : 60**

**Note :** This paper is of **Sixty (60)** marks containing **three (03)** sections A, B and C. Learners are required to attempt the questions contained in these sections according to the detailed instructions given therein.

#### Section-A

##### (Long Answer Type Questions)

**Note :** Section 'A' contains four (04) long answer type questions of fifteen (15) marks each. Learners are required to answer *two* (02) questions only.

1. Reduce the differential equation :

$$\phi''(x) - 3\phi'(x) + 2\phi(x) = 4 \sin x$$

with the conditions

$\phi(0) = 1$ ,  $\phi'(0) = -2$ , into a non-homogenous Volterra's integral equation of second kind.

2. Solve  $(D^2 + 9)y = \cos 2t$  if  $y(0) = 1$ ,  $y\left(\frac{\pi}{2}\right) = -1$

and  $D \equiv \frac{d}{dx}$ .

- Find the Fourier sine and cosine transform of  $e^{-x}$  and using the inversion formulae, recover the original functions in both cases.
- If  $f^{(n)}(x)$  is the  $n$ th derivative of  $f(x)$  with respect to  $x$ , then prove that :

$$M[f^{(n)}(x); P] = \frac{(-1)^n \sqrt{P}}{\sqrt{P-n}} M[f(P-n)]$$

### Section-B

#### (Short Answer Type Questions)

**Note :** Section 'B' contains eight (08) short answer type questions of five (05) marks each. Learners are required to answer *four* (04) questions only.

- Prove that :  $\phi(x) = (1+x^2)^{-3/2}$  is a solution of Volterra integral equation :

$$\phi(x) = \frac{1}{1+x^2} - \int_0^x \left( \frac{t}{1+t^2} \right) \phi(t) dt$$

- Form an integral equation corresponding to the differential equation :

$$\frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = 0,$$

with the initial conditions  $y(0) = 1$ ,  $y'(0) = 0$ .

- Find the resolvent kernel of integral equation  $\phi''(x) - 2\phi'(x) + \phi(x) = 0$ ;  $\phi(t) = 0$ ,  $\phi'(t) = 1$  if  $k(x, t) = 2 - (x - t)$ ,  $\lambda = 1$ .
- State and prove second shifting theorem for Laplace transform.

5. For Mellin transform, prove that :

$$M\left[\frac{1}{x} f\left(\frac{1}{x}\right); P\right] = M f(1 - P)$$

6. Find the Hankel transform of  $\frac{df}{dx}$  when  $f = \frac{e^{-ax}}{x}$  and

$$n = 1.$$

7. Find :

$$L^{-1}\left\{\frac{e^{4-3P}}{(P+4)^{5/2}}\right\}$$

8. Evaluate :

$$\int_0^\infty \left(\frac{e^{-at} - e^{-bt}}{t}\right) dt$$

### Section-C

#### (Objective Type Questions)

**Note :** Section 'C' contains ten (10) objective type questions of one (01) mark each. All the questions of this section are compulsory.

1. An integral equation in Fredholm type if :

- (i) Both the limits are constants
- (ii) Both the limits are variable
- (iii) Only upper limit is variable
- (iv) Only lower limit is variable

2. In integral equations,  $G(x, t)$  is called :

- (i) Fredholm's function

- (ii) Green's function
  - (iii) Volterra's function
  - (iv) None of these
3. Equation, which and unknown function appears under the integral sign is called as :
- (i) Linear differential equation
  - (ii) Partial integral equation
  - (iii) Linear integral equation
  - (iv) Gaussian integral equation
4. The solution, corresponding to eigen values of  $\lambda$ , can be expressed as :
- (i) Sum of eigen functions
  - (ii) Arbitrary multiples of eigen functions
  - (iii) Difference of eigen functions
  - (iv) None of these
5.  $L \{ \sinh at \}$  is,  $P > |a|$  :
- (i)  $\frac{a}{p^2 + a^2}$
  - (ii)  $\frac{P}{p^2 + a^2}$
  - (iii)  $\frac{P}{p^2 - a^2}$
  - (iv) None of these

6. A kernel  $K(x, t)$  of Fredholm integral equation is said to be ..... if it can be expressed as the sum of a finite number of terms, each of which is the product of a function of  $x$  alone and a function of  $t$  alone :

- (i) Non-degenerate
- (ii) Separable
- (iii) Non-separable
- (iv) Degenerate

7. Hankel transform of  $\left\{ \frac{e^{-x}}{x} \right\}$  is :

- (i)  $(1 + P^2)^{-1/2}$
- (ii)  $(1 + P)^{-1/2}$
- (iii)  $(1 + P^2)^{-3/2}$
- (iv) None of these

8. In Mellin transform,  $M[x^a f(x); P] = :$

- (i)  $M f(x - a)$
- (ii)  $M f(x / a)$
- (iii)  $M f(x + a)$
- (iv) None of these

9.  $L\{t^2 e^{2t}\}$  is :

- (i)  $\frac{2}{(P - 2)^2}$

$$(ii) \quad \frac{P}{(P-2)^3}$$

$$(iii) \quad \frac{2}{(P-2)^3}$$

$$(iv) \quad \frac{2}{(P+2)^3}$$

10.  $L\left(\frac{e^{at}-1}{a}\right)$  is,  $P > 0$  :

$$(i) \quad \frac{1}{P(P+a)}$$

$$(ii) \quad \frac{1}{P(P-a)}$$

$$(iii) \quad \frac{P}{P-a}$$

(iv) None of these